

CHERN NUMBERS AND DIFFEOMORPHISM TYPES OF PROJECTIVE VARIETIES

D. KOTSCHICK

Herrn Prof. Dr. F. Hirzebruch zum 80. Geburtstag gewidmet.

ABSTRACT. In 1954 Hirzebruch asked which linear combinations of Chern numbers are topological invariants of smooth complex projective varieties. We give a complete answer to this question in small dimensions, and also prove partial results without restrictions on the dimension.

1. INTRODUCTION

More than fifty years ago, Hirzebruch raised the question to what extent the Chern and Hodge numbers of projective algebraic manifolds are topologically invariant, see Problem 31 in [2]. He noted that Chern numbers of almost complex manifolds are not topologically invariant simply because there are too many almost complex structures, even on a fixed manifold. Then, in 1959, Borel and Hirzebruch gave an example of a 10-dimensional closed oriented manifold with two projective algebraic structures for which c_1^5 are different, see [1] Section 24.11. Until 1987, when the commentary in [4] was written, nothing further was discovered concerning this question. In particular, Hirzebruch wrote then that he did not know whether c_1^3 and c_1^4 are topological invariants of complex three- and four-folds respectively.

In this paper we prove that in complex dimension 3 the only linear combinations of the Chern numbers c_1^3 , c_1c_2 and c_3 that are invariant under orientation-preserving diffeomorphisms of simply connected projective algebraic manifolds are the multiples of the Euler characteristic c_3 . In dimension 4 the only linear combinations of Chern numbers that are invariant are the linear combinations of the Euler characteristic c_4 and of the Pontryagin numbers p_1^2 and p_2 . We also prove some partial results in arbitrary dimensions.

These results stem from the fact that in complex dimension 2 the Chern number c_1^2 is not invariant under orientation-reversing homeomorphisms; cf. [5]. By suitable stabilisation of the counterexamples, we find enough examples at least in dimensions 3 and 4 to detect the independent variation of all Chern numbers which are not combinations of the Euler and Pontryagin numbers. Our results suggest a weaker form of Hirzebruch's problem, asking whether the topology determines the Chern numbers up to finite ambiguity. We have no counterexample to an affirmative answer to this weaker question in the projective algebraic case, although it is known to be false for non-Kähler complex manifolds; cf. [7].

2. PRELIMINARY RESULTS

2.1. Complex surfaces. For complex surfaces there are two Chern numbers, c_2 and c_1^2 , which turn out to be diffeomorphism invariants even without assuming that the diffeomorphism is orientation-preserving with respect to the orientations given by the complex structures:

Date: December 20, 2007; MSC 2000: primary 57R20, secondary 14J30, 14J35, 32Q55.

Theorem 1. *If two compact complex surfaces are diffeomorphic, then their Chern numbers coincide.*

Proof. In this case c_2 is the topological Euler characteristic e . By Wu's formula we have

$$(1) \quad c_1^2(X) = 2e(X) + p_1(X) .$$

The first Pontryagin number is 3 times the signature, and so the right-hand side of (1) is invariant not just under orientation-preserving diffeomorphisms, but even under orientation-preserving homotopy equivalences¹.

Now suppose that two compact complex surfaces are orientation-reversing diffeomorphic, with respect to the orientations defined by their complex structures. Then, using Seiberg–Witten theory, I proved in 1995, see Theorem 2 of [6], that the signatures of these surfaces vanish. Thus, their Chern numbers agree by (1). $\square$

The statement about orientation-reversing diffeomorphisms concerns projective algebraic surfaces only, because, by the classification of complex surfaces, a complex surface with positive signature is always projective.

As we saw in the proof, c_1^2 is invariant under orientation-preserving homotopy equivalences, and under orientation-reversing diffeomorphisms. But it is not invariant under orientation-reversing homeomorphisms. Already in 1991, I had proved the following:

Theorem 2 ([5]). *There are infinitely many pairs of simply connected projective algebraic surfaces X_i and Y_i of non-zero signature which are orientation-reversing homeomorphic.*

The proof is based on geography results for surfaces of general type due to Persson and Chen. The surfaces X_i and Y_i are projective algebraic because they are of general type. They can be chosen to contain embedded holomorphic spheres, in which case they can not be orientation-reversing diffeomorphic, although they are orientation-reversing homeomorphic. This was the motivation for the results of [6] quoted in the proof of Theorem 1.

By Wu's formula (1) the homeomorphic surfaces X_i and Y_i have different c_1^2 . Indeed, the homeomorphism in question preserves the Euler number and reverses the sign of the signature, so that (1) gives:

$$(2) \quad c_1^2(Y_i) = 4e(X_i) - c_1^2(X_i) .$$

Wu's formula (1) shows in particular that the unoriented homeomorphism type almost determines the Chern numbers of a compact complex surface: there are only two possible values for c_1^2 (and only one for c_2 , of course).

We shall use the examples from Theorem 2 as building blocks for our high-dimensional examples.

2.2. Inductive formulae for Chern classes. We require the following easy calculation.

Lemma 1. *Let A be a compact complex n -fold, and $B = A \times \mathbb{C}P^1$. Then the Chern numbers of B are*

$$(3) \quad c_{r_1} \dots c_{r_k}(B) = 2 \sum_{j=1}^k c_{r_1} \dots c_{r_{j-1}} \dots c_{r_k}(A) .$$

¹All this was known in 1954, and Hirzebruch [2] remarked that the Chern numbers of an algebraic surface are topological invariants (of the underlying oriented manifold).

Proof. The Whitney sum formula $c(TB) = c(TA)c(T\mathbb{C}P^1)$ for the total Chern classes implies that, with respect to the Künneth decomposition of the cohomology of B , the Chern classes of B are

$$\begin{aligned} c_1(B) &= c_1(A) + c_1(\mathbb{C}P^1) \\ c_2(B) &= c_2(A) + c_1(A)c_1(\mathbb{C}P^1) \\ &\dots \\ c_n(B) &= c_n(A) + c_{n-1}(A)c_1(\mathbb{C}P^1) \\ c_{n+1}(B) &= c_n(A)c_1(\mathbb{C}P^1) . \end{aligned}$$

The claim follows using that the first Chern number of $\mathbb{C}P^1$ equals 2. $\square$

We also need the following generalization of Lemma 1 to non-trivial $\mathbb{C}P^1$ -bundles:

Lemma 2. *Let B be a compact complex surface and $E \rightarrow B$ a holomorphic vector bundle of rank two. Then the projectivisation $X = \mathbb{P}(E)$ has*

$$(4) \quad \begin{aligned} c_3(X) &= 2c_2(B) , \\ c_1c_2(X) &= 2(c_1^2(B) + c_2(B)) , \\ c_1^3(X) &= 6c_1^2(B) + 2p_1(\mathbb{P}(E)) . \end{aligned}$$

Here $p_1(\mathbb{P}(E)) = c_1^2(E) - 4c_2(E)$ is the first Pontryagin number for the group $SO(3) = PU(2)$, which is the structure group of the sphere bundle $X \rightarrow B$. Notice that in the case that $p_1(\mathbb{P}(E)) = 0$, the formulae reduce to those obtained for the trivial bundle.

Proof. The formulae for c_3 and for c_1c_2 are immediate from the multiplicativity of the topological Euler characteristic and of the Todd genus, recalling that the Todd genera in dimension 2 and 3 are $\frac{1}{12}(c_1^2 + c_2)$, respectively $\frac{1}{24}c_1c_2$. To compute c_1^3 note that by the Leray-Hirsch theorem the cohomology ring of X is generated as a $H^*(B)$ -module by a class $y \in H^2(X)$ restricting as a generator to every fiber and satisfying the relation

$$y^2 + c_1(E)y + c_2(E) = 0 .$$

Moreover, $c_1(X) = c_1(B) + c_1(E) + 2y$ because the vertical tangent bundle has first Chern class $c_1(E) + 2y$. The third power is computed straightforwardly using the relation and the fact that y evaluates to 1 on the fiber. $\square$

3. COMPLEX THREE-FOLDS

A variant of Hirzebruch's problem for three-folds was taken up by LeBrun in 1998, see [7], who proved that there are closed 6-manifolds which admit complex structures with different c_1c_2 and c_1^3 . He even proved that a fixed manifold can have complex structures realising infinitely many different values for c_1c_2 . However, for all the examples discussed in [7] only one of the complex structures is projective algebraic, or at least Kähler, and all the others are non-Kähler. Therefore, these examples say nothing about the topological invariance of Chern numbers for projective algebraic three-folds.

Nevertheless, both c_1c_2 and c_1^3 are not diffeomorphism invariants of projective three-folds:

Proposition 1. *There are infinitely many pairs of projective algebraic three-folds Z_i and T_i with the following properties:*

- (i) Each Z_i and T_i admits an orientation-reversing diffeomorphism.
- (ii) For each i the manifolds underlying Z_i and T_i are diffeomorphic.
- (iii) For each i one has $c_1c_2(Z_i) \neq c_1c_2(T_i)$ and $c_1^3(Z_i) \neq c_1^3(T_i)$.

Proof. Let X_i and Y_i be the algebraic surfaces from Theorem 2, constructed in [5], and take $Z_i = X_i \times \mathbb{C}P^1$ and $T_i = Y_i \times \mathbb{C}P^1$. Then the identity on the first factor times complex conjugation on the second factor gives an orientation-reversing selfdiffeomorphism of Z_i and of T_i .

Denote by $\bar{Y}_i$ the smooth manifold underlying Y_i , but endowed with the orientation opposite to the one induced by the complex structure. Then X_i and $\bar{Y}_i$ are orientation-preserving homeomorphic simply connected smooth four-manifolds, and are therefore h-cobordant. If W is an h-cobordism between them, then $W \times S^2$ is an h-cobordism between Z_i and $\bar{T}_i = \bar{Y}_i \times \mathbb{C}P^1$. By Smale's h-cobordism theorem, Z_i and $\bar{T}_i$ are orientation-preserving diffeomorphic. As Z_i and T_i admit orientation-reversing diffeomorphisms, we conclude that they are both orientation-preserving and orientation-reversing diffeomorphic.

For the Chern numbers (3) gives

$$(5) \quad \begin{aligned} c_1c_2(Z_i) &= 2(c_1^2 + c_2)(X_i), \\ c_1^3(Z_i) &= 6c_1^2(X_i), \end{aligned}$$

and similarly for T_i and Y_i . As X_i and Y_i have the same c_2 but different c_1^2 , we conclude that Z_i and T_i have different c_1c_2 and different c_1^3 . $\square$

Thus c_1c_2 and c_1^3 are not topological invariants of projective three-folds, but it is not yet clear that they vary independently. This is the content of the following:

Theorem 3. *The only linear combinations of the Chern numbers c_1^3 , c_1c_2 and c_3 that are invariant under orientation-preserving diffeomorphisms of simply connected projective algebraic three-folds are the multiples of the Euler characteristic c_3 .*

Proof. First of all, let us dispose of the orientation question. If two complex three-folds are orientation-reversing diffeomorphic with respect to the orientations given by their complex structures, then they become orientation-preserving diffeomorphic after we replace one of the complex structures by its complex conjugate. As the conjugate complex structure has the same Chern numbers as the original one, we do not have to distinguish between orientation-preserving and orientation-reversing diffeomorphisms.

All the examples constructed in the proof of Proposition 1 have the property that $3c_1c_2 - c_1^3$ agrees on Z_i and T_i , as follows by combining (2) with (5) and the topological invariance of c_2 for surfaces. But, by Proposition 1, linear combinations of $3c_1c_2 - c_1^3$ and of c_3 are the only candidates left for combinations of Chern numbers that can be topological invariants of projective three-folds. In order to show that $3c_1c_2 - c_1^3$ is not an oriented diffeomorphism invariant we shall use certain ruled manifolds which are non-trivial $\mathbb{C}P^1$ -bundles, rather than the products used above.

Consider again a pair X_i and Y_i of simply connected algebraic surfaces as in Theorem 2. For simplicity we just denote them by X and Y , with orientations implicitly given by the complex structures. The oriented manifolds X and $\bar{Y}$ are orientation-preserving h-cobordant. Let M be the projectivisation of the holomorphic tangent bundle TY of Y . Temporarily ignoring the complex structure of M , we think of it as a smooth oriented two-sphere bundle over Y , or over $\bar{Y}$. If W is any h-cobordism between X and $\bar{Y}$, then the two-sphere bundle $M \rightarrow \bar{Y}$ extends to a uniquely defined oriented two-sphere bundle $V \rightarrow W$. Let N be the restriction of this bundle to $X \subset W$. If we give N the orientation induced from that of X and M the orientation induced from that of $\bar{Y}$,

then, by construction, V is an h-cobordism between N and M . By Smale's h-cobordism theorem, M and N are diffeomorphic.

Because the bundle $p: M \rightarrow Y$ was defined as the projectivisation of the holomorphic tangent bundle of Y , its characteristic classes are $w_2(p) = w_2(Y)$ and $p_1(p) = c_1^2(Y) - 4c_2(Y)$. Considered as a bundle over $\bar{Y}$, p has the same Stiefel–Whitney class, but the first Pontryagin number changes sign. It follows that $q: N \rightarrow X$ has

$$p_1(q) = -c_1^2(Y) + 4c_2(Y) = c_1^2(X) ,$$

where the last equality is from (2). Moreover, $w_2(q) = w_2(X)$, although X and $\bar{Y}$ are not diffeomorphic. This follows for example from the cohomological characterisation of $w_2(X)$ as the unique element of $H^2(X; \mathbb{Z}_2)$ which for all x satisfies

$$w_2(X) \cdot x \equiv x^2 \pmod{2} .$$

The bundle q is determined by $w_2(q) = w_2(X)$ and $p_1(q) = c_1^2(X)$, and so we can think of it as the projectivisation of the holomorphic rank two bundle $\mathcal{O}(K) \oplus \mathcal{O} \rightarrow X$. Therefore the total space N inherits a complex-algebraic structure from that of X . Its Chern numbers are given by (4):

$$\begin{aligned} c_3(N) &= 2c_2(X) , \\ (6) \quad c_1c_2(N) &= 2(c_1^2(X) + c_2(X)) , \\ c_1^3(N) &= 8c_1^2(X) . \end{aligned}$$

This N is diffeomorphic to M , which has a complex-algebraic structure as the projectivisation of the holomorphic tangent bundle of Y . (Recall from the beginning of the proof that we do not have to keep track of the orientations induced by complex structures, because we can always replace a structure by its complex conjugate.) The Chern numbers of M are also given by (4):

$$\begin{aligned} c_3(M) &= 2c_2(Y) = 2c_2(X) , \\ (7) \quad c_1c_2(M) &= 2(c_1^2(Y) + c_2(Y)) = 2(-c_1^2(X) + 5c_2(X)) , \\ c_1^3(M) &= 8c_1^2(Y) - 8c_2(Y) = 8(-c_1^2(X) + 3c_2(X)) , \end{aligned}$$

using (2) to replace the Chern numbers of Y by combinations of those of X . Unlike for the examples in Proposition 1, the combination $3c_1c_2 - c_1^3$ is not the same for M and N . This finally shows that c_1c_2 and c_1^3 vary independently (within a fixed diffeomorphism type). $\square$

Although the Chern numbers of a projective three-fold are not determined by the underlying differentiable manifold, this may still be the case up to finite ambiguity. By the Hirzebruch–Riemann–Roch theorem one has

$$(8) \quad \frac{1}{24}c_1c_2 = 1 - h^{1,0} + h^{2,0} - h^{3,0} ,$$

so that in the Kähler case c_1c_2 is bounded from above and from below by linear combinations of Betti numbers. In particular, for Kähler structures on a fixed 6-manifold c_1c_2 can take at most finitely many values. We are left with the following:

Problem 1. *Does c_1^3 take on only finitely many values on the projective algebraic structures with the same underlying 6-manifold?*

The issue here is that there is no Riemann–Roch type formula expressing c_1^3 as a combination of Hodge numbers and the other Chern numbers.

For three-folds with ample canonical bundle one has $c_1^3 < 0$, and Yau's celebrated work [11] gives $c_1^3 \geq \frac{8}{3}c_1c_2$. As c_1c_2 is bounded below by a linear combination of Betti numbers, we have a positive answer to Problem 1 for this restricted class of projective three-folds. Even in the non-Kähler category, there are no examples where infinitely many values are known to arise for c_1^3 .

4. HIGHER DIMENSIONS

It is now very easy to show that, except for the Euler number, no Chern number is diffeomorphism-invariant:

Theorem 4. *For projective algebraic n -folds with $n \geq 3$ the only Chern number c_I which is diffeomorphism-invariant is the Euler number c_n .*

Note that by Theorem 1 this is false for $n = 2$, because in that case c_1^2 is also diffeomorphism-invariant. On the other hand, by Theorem 2 it is not homeomorphism-invariant, so that Theorem 4 is true for $n = 2$ if we replace diffeomorphism-invariance by homeomorphism-invariance. As in the case of Proposition 1, the examples we exhibit in the proof of Theorem 4 admit orientation-reversing diffeomorphisms, so that one cannot restore diffeomorphism-invariance of $c_I \neq c_n$ by restricting to orientation-preserving diffeomorphisms only.

Proof. For $n = 3$ this was already proved in Proposition 1. For $n > 3$ we take the examples T_i and Z_i from Proposition 1 and multiply them by $n - 3$ copies of $\mathbb{C}P^1$. Call these products T'_i and Z'_i . Using formula (3) and induction, we see that, on the one hand, c_n is a universal multiple of the c_2 of the surfaces we started with. On the other hand, $c_1^n(T'_i)$ and $c_1^n(Z'_i)$ are universal multiples of $c_1^2(X_i)$ and of $c_1^2(Y_i)$ respectively, and so are different. All other Chern numbers c_I are universal linear combinations of $c_2(X_i)$ and $c_1^2(X_i)$, respectively $c_2(Y_i)$ and $c_1^2(Y_i)$, with the coefficients of both c_2 and c_1^2 strictly positive. As X_i and Y_i have the same c_2 but different c_1^2 , the result follows. $\square$

Although the individual Chern numbers are not diffeomorphism-invariant, certain linear combinations are invariant once we restrict to orientation-preserving diffeomorphisms. Of course, as remarked by Hirzebruch [4], the Pontryagin numbers p_J have this invariance property², but this only helps when the complex dimension is even.

Problem 2. *Prove that, in arbitrary dimensions, the only combinations of Chern numbers that are invariant under orientation-preserving diffeomorphisms of smooth complex projective varieties are linear combinations of Euler and Pontryagin numbers.*

For complex dimension 3 this is Theorem 3 above. Theorem 5 below deals with the case of complex dimension 4.

It would be interesting to know whether each of the Chern numbers $c_I \neq c_n$ takes on only finitely many values on a fixed smooth manifold. In the Kähler case this is known to be true for c_1c_{n-1} by a result of Libgober and Wood, who showed that this Chern number is always a linear combination of Hodge numbers, see Theorem 3 in [8]. In the non-Kähler case c_1c_{n-1} can take on infinitely many values on a fixed manifold. This follows as in the proof of Theorem 4 by taking products of LeBrun's examples [7] mentioned in the previous section with $\mathbb{C}P^1$, and using formula (3). Because c_1c_2 takes on infinitely many values on a fixed 6-manifold, the same conclusion holds for c_1c_{n-1} in real dimension $2n \geq 8$.

²Note that, unlike the Euler number, the Pontryagin numbers change sign under a change of orientation.

Returning to the Kähler case, the Riemann–Roch theorem expresses the χ_p -genus³

$$\chi_p = \sum_{q=0}^n (-1)^q h^{p,q}$$

as a linear combination of Chern numbers, and it follows that the combinations of Chern numbers which appear in this way can take on only finitely many values on a fixed manifold, as they are bounded above and below by linear combinations of Betti numbers.

Remark 1. If the complex dimension n is odd, then the Todd genus expressing the Euler characteristic $\chi_0 = (-1)^n \chi_n$ of the structure sheaf as a combination of Chern numbers does not involve c_1^n . This follows from the *Bemerkungen* in Section 1.7 of [3]. On the one hand, in any dimension, the coefficient of c_n in the Todd genus agrees with the coefficient of c_1^n . On the other hand, the Todd genus is divisible by c_1 if n is odd.

Generalising this Remark, and what we saw for $n = 3$ in the previous section, we now prove:

Proposition 2. *If M is Kähler of odd complex dimension $n > 1$, then all χ_p are linear combinations of Chern numbers which do not involve c_1^n .*

This shows that for odd n there is no general way to extract the value of c_1^n from the Hodge numbers. In particular, one can not obtain a finiteness result for c_1^n in this way.

Proof. The Kähler symmetries imply $\chi_p = (-1)^n \chi_{n-p}$, so that it is enough to prove the claim for $p > \frac{n}{2}$. We shall do this by descending induction starting at $p = n$.

Salamon [10] proved that for $2 \leq k \leq n$ the number

$$(9) \quad \sum_{p=k}^n (-1)^p \binom{p}{k} \chi_p$$

is a linear combination of Chern numbers each of which involves a c_i with $i > n - 2[\frac{k}{2}]$, see [10] Corollary 3.3. Using this for n odd and $k = n$, we obtain once more the claim for χ_n treated already in the Remark above.

Suppose now that the claim has been proved for $\chi_n, \chi_{n-1}, \dots, \chi_j$ with $j > \frac{n}{2} + 1$. Then we consider (9) with $k = j - 1$. (Note that this still satisfies $k \geq 2$.) As χ_p with $p \geq j$ does not involve c_1^n by the induction hypothesis, Salamon's result implies that χ_{j-1} does not involve c_1^n either. $\square$

5. FOUR-FOLDS

In the case of four-folds, in addition to the Euler number c_4 , the following are invariants of the underlying oriented smooth manifold:

$$(10) \quad \begin{aligned} p_1^2 &= (c_1^2 - 2c_2)^2 = c_1^4 - 4c_1^2 c_2 + 4c_2^2 \\ p_2 &= c_2^2 - 2c_1 c_3 + 2c_4. \end{aligned}$$

The vector space of Chern numbers of four-folds is 5-dimensional, containing the 3-dimensional subspace spanned by c_4, p_1^2 and p_2 . It turns out that all combinations of Chern numbers that are invariant under orientation-preserving diffeomorphisms are contained in this subspace:

³Our notation is consistent with [10], changing the traditional superscript in χ^p from [3] to a subscript.

Theorem 5. *The only linear combinations of Chern numbers that are invariant under orientation-preserving diffeomorphisms of simply connected projective algebraic four-folds are linear combinations of the Euler characteristic and of the Pontryagin numbers.*

Proof. This is a rather formal consequence of our results for complex three-folds. Consider the vector space of Chern number triples (c_3, c_2c_1, c_1^3) . Whenever we have a smooth six-manifold with two different complex structures, the difference of the two Chern vectors must be in the kernel of any linear functional corresponding to a topologically invariant combination of Chern numbers. In the proof of Theorem 3 we produced two kinds of examples for which these difference vectors were linearly independent. Therefore the space of topologically invariant combinations of Chern numbers is at most one-dimensional, and as it contains c_3 it is precisely one-dimensional.

Consider now the four-folds obtained by multiplying the three-dimensional examples by $\mathbb{C}P^1$. If the difference of Chern vectors in a three-dimensional example is $(0, a, b)$, then by (3) the difference of Chern vectors $(c_4, c_1c_3, c_2^2, c_1^2c_2, c_1^4)$ for the product with $\mathbb{C}P^1$ is $(0, 2a, 4a, 4a + 2b, 8b)$. Two examples in dimension three with linearly independent difference vectors lead to examples in dimension four which also have linearly independent difference vectors. Thus, in the five-dimensional space spanned by the Chern numbers of complex projective four-folds, the subspace invariant under orientation-preserving diffeomorphisms has codimension at least two. As it contains the linearly independent elements c_4, p_1^2 and p_2 , it is exactly three-dimensional. $\square$

Concerning the weaker question which Chern numbers of projective or Kähler four-folds are determined by the topology up to finite ambiguity, this is so for c_1c_3 on general grounds, see the discussion above and [8, 10]. The formula for p_2 then shows that c_2^2 is also determined up to finite ambiguity. Using either the formula for p_1^2 or the Riemann–Roch formula for the structure sheaf, we conclude that $c_1^4 - 4c_1^2c_2$ is also determined up to finite ambiguity, but it is not clear whether this is true for c_1^4 and $c_1^2c_2$ individually. Note that a negative answer to Problem 1, giving infinitely many values for c_1^3 on a fixed 6-manifold, would show that c_1^4 also takes on infinitely many values on a fixed 8-manifold by taking products with $\mathbb{C}P^1$.

For four-folds with ample canonical bundle one has $c_1^4 > 0$, and Yau’s work [11] gives $c_1^4 \leq \frac{5}{2}c_1^2c_2$. Therefore

$$0 < c_1^4 \leq \frac{5}{3}(4c_1^2c_2 - c_1^4) .$$

As the right-hand side takes on only finitely many values, the same is true for c_1^4 , and then for $c_1^2c_2$ as well.

Remark 2. Pasquotto [9] recently raised the question of the topological invariance of Chern numbers of symplectic manifolds, particularly in (real) dimensions 6 and 8. Our results for Kähler manifolds of course show that Chern numbers of symplectic manifolds are not topological invariants. In the Kähler case we have used Hodge theory to argue that the variation of Chern numbers is quite restricted, often to finitely many possibilities. It would be interesting to know whether any finiteness results hold in the symplectic non-Kähler category.

REFERENCES

1. A. Borel and F. Hirzebruch, *Characteristic classes and homogeneous spaces, II*, Amer. J. Math. **81** (1959), 315–382.
2. F. Hirzebruch, *Some problems on differentiable and complex manifolds*, Ann. Math. **60** (1954), 213–236.
3. F. Hirzebruch, *Neue topologische Methoden in der algebraischen Geometrie*, 2. ergänzte Auflage, Springer Verlag 1962.

4. F. Hirzebruch, *Gesammelte Abhandlungen*, Band I, Springer-Verlag 1987.
5. D. Kotschick, *Orientation-reversing homeomorphisms in surface geography*, Math. Annalen **292** (1992), 375–381.
6. D. Kotschick, *Orientations and geometrisations of compact complex surfaces*, Bull. London Math. Soc. **29** (1997), 145–149.
7. C. LeBrun, *Topology versus Chern numbers for complex 3-folds*, Pacific J. Math. **191** (1999), 123–131.
8. A. S. Libgober and J. W. Wood, *Uniqueness of the complex structure on Kähler manifolds of certain homotopy types*, J. Differential Geometry **32** (1990), 139–154.
9. F. Pasquotto, *Symplectic geography in dimension 8*, Manuscripta math. **116** (2005), 341–355.
10. S. M. Salamon, *On the cohomology of Kähler and hyper-Kähler manifolds*, Topology **35** (1996), 137–155.
11. S.-T. Yau, *Calabi’s conjecture and some new results in algebraic geometry*, Proc. Natl. Acad. Sci. USA **74** (1977), 1798–1799.

MATHEMATISCHES INSTITUT, LUDWIG-MAXIMILIANS-UNIVERSITÄT MÜNCHEN, THERESIENSTR. 39, 80333 MÜNCHEN, GERMANY

E-mail address: dieter@member.ams.org